

If you are this age, talk to your healthcare professional about these vaccines

If you are this age, ↓	Flu <i>Influenza</i>	Td/Tdap Tetanus, diphtheria, pertussis	Shingles <i>Zoster</i>	Pneumococcal		Meningococcal		MMR Measles, mumps, rubella	HPV <i>Human papillomavirus</i>		Chickenpox <i>Varicella</i>	Hepatitis A	Hepatitis B	Hib <i>Haemophilus influenzae</i> type b
				PCV13	PPSV23	MenACWY or MPSV4	MenB		for women	for men				
19 - 21 years	Green	Green	Light Blue	Blue	Blue	Blue	Blue	Green	Green	Green	Green	Blue	Blue	Blue
22 - 26 years	Green	Green	Light Blue	Blue	Blue	Blue	Blue	Green	Green	Blue	Green	Blue	Blue	Blue
27 - 59 years	Green	Green	Light Blue	Blue	Blue	Blue	Blue	Green	Light Blue	Light Blue	Green	Blue	Blue	Blue
60 - 64 years	Green	Green	Green	Blue	Blue	Blue	Blue	Light Blue	Light Blue	Light Blue	Green	Blue	Blue	Blue
65+ year	Green	Green	Green	Blue	Blue	Blue	Blue	Light Blue	Light Blue	Light Blue	Green	Blue	Blue	Blue

More Information:

You should get flu vaccine every year.

You should get a Td booster every 10 years. You also need 1 dose of Tdap. Women should get a Tdap vaccine during every pregnancy to help protect the baby.

You should get shingles vaccine even if you have had shingles before.

You should get 1 dose of PCV13 and at least 1 dose of PPSV23 depending on your age and health condition.

You should get this vaccine if you did not get it when you were a child.

You should get HPV vaccine if you are a woman through age 26 years or a man through age 21 years and did not already complete the series.



Recommended For You: This vaccine is recommended for you *unless* your healthcare professional tells you that you do not need it or should not get it.



May Be Recommended For You: This vaccine is recommended for you if you have certain risk factors due to your health condition or other. Talk to your healthcare professional to see if you need this vaccine.

If you are traveling outside the United States, you may need additional vaccines.

Ask your healthcare professional about which vaccines you may need at least 6 weeks before you travel.

For more information, call 1-800-CDC-INFO (1-800-232-4636) or visit www.cdc.gov/vaccines



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

If you have this health condition,

talk to your healthcare professional about these vaccines

	Flu <i>Influenza</i>	Td/Tdap Tetanus, diphtheria, pertussis	Shingles Zoster	Pneumococcal		Meningococcal		MMR Measles, mumps, rubella	HPV <i>Human papillomavirus</i>		Chickenpox <i>Varicella</i>	Hepatitis A	Hepatitis B	Hib <i>Haemophilus influenzae</i> type b
				PCV13	PPSV23	MenACWY or MPSV4	MenB		for women	for men				
Pregnancy	Green	Green	Light Purple	Green	Green	Blue	Blue	Light Purple	Green	Green	Light Purple	Blue	Blue	Green
Weakened Immune System	Green	Green	Light Purple	Green	Green	Blue	Blue	Light Purple	Green	Green	Light Purple	Blue	Blue	Green
HIV: CD4 count less than 200	Green	Green	Light Purple	Green	Green	Blue	Blue	Light Purple	Green	Green	Light Purple	Blue	Blue	Green
HIV: CD4 count 200 or greater	Green	Green	Green	Green	Green	Blue	Blue	Green	Green	Green	Green	Blue	Blue	Green
Kidney disease or poor kidney function	Green	Green	Green	Green	Green	Blue	Blue	Green	Green	Green	Green	Blue	Blue	Green
Asplenia (if you do not have a spleen or if it does not work well)	Green	Green	Green	Green	Green	Blue	Blue	Green	Green	Green	Green	Blue	Blue	Green
Heart disease Chronic lung disease Chronic alcoholism	Green	Green	Green	Green	Green	Blue	Blue	Green	Green	Green	Green	Blue	Blue	Green
Diabetes (Type 1 or Type 2)	Green	Green	Green	Green	Green	Blue	Blue	Green	Green	Green	Green	Blue	Blue	Green
Chronic Liver Disease	Green	Green	Green	Green	Green	Blue	Blue	Green	Green	Green	Green	Blue	Blue	Green

More Information:

You should get flu vaccine every year.

You should get a Td booster every 10 years. You also need 1 dose of Tdap vaccine. Women should get Tdap vaccine during every pregnancy.

You should get shingles vaccine if you are age 60 years or older, even if you have had shingles before.

You should get 1 dose of PCV13 and at least 1 dose of PPSV23 depending on your age and health condition.

You should get this vaccine if you did not get it when you were a child.

You should get HPV vaccine if you are a woman through age 26 years or a man through age 21 years and did not already complete the series.

You should get Hib vaccine if you do not have a spleen, have sickle cell disease, or received a bone marrow transplant.

Recommended For You: This vaccine is recommended for you *unless* your healthcare professional tells you that you do not need it or should not get it.

May Be Recommended For You: This vaccine is recommended for you if you have certain other risk factors due to your age, health condition or other. Talk to your healthcare professional to see if you need this vaccine.

YOU SHOULD NOT GET THIS VACCINE



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

For more information, call 1-800-CDC-INFO (1-800-232-4636) or visit www.cdc.gov/vaccines