



Sanitizer Log



Sanitizer solution shall be maintained clean and at proper concentration of 50-200 ppm of chlorine (bleach) or 150-400 ppm quaternary ammonium.

Sanitizer shall be mixed with lukewarm water 75°F.

Use test strips appropriate for the sanitizer to ensure proper concentration (i.e. white strips for chlorine).

Sanitizer buckets shall be changed as needed or a minimum of every 4 hours.

| Date | Time | Sanitizer Buckets or Spray Bottles 50-200 ppm chlorine or 150-400 ppm quaternary ammonium | Sanitizer Dish Machines/ Sink 50-200 ppm chlorine or Rinse temperature a minimum of 180°F. | Date | Time | Sanitizer Buckets or Spray Bottles 50-200 ppm chlorine or 150-400 ppm quaternary ammonium | Sanitizer Dish Machines/ Sink 50-200 ppm chlorine or Rinse temperature a minimum of 180°F. |
|------|------|--|--|------|------|--|--|
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |



HELP PREVENT FOOD-RELATED ILLNESS